

- GRIGORE, M. T. & J. WINDIS. 1994. Decline of the Karner blue butterfly in the Oak Openings of northwest Ohio, pp 135–142. *In* D. A. Andow, R. J. Baker & C. P. Lane (eds.), Karner blue butterfly: symbol of a vanishing landscape. Misc. Publ. 84-1994, Minn. Agric. Exp. Stat., Univ. Minn. St. Paul.
- HAFERNIK, J. E., JR. 1992. Threats to invertebrate biodiversity: implications for conservation strategies, pp. 171–195. *In* P. L. Fiedler & S. K. Jain (eds.), Conservation biology: the theory and practice of nature conservation, preservation, and management. Chapman and Hall, New York.
- IFTNER, D. C., J. A. SHUEY & J. V. CALHOUN. 1992. Butterflies and skippers of Ohio. Ohio Biol. Surv. Bull. New Series Vol. 9, No. 1. 212 pp.
- METZLER, E. H. & V. P. LUCAS. 1990. An endangered moth in Ohio, with notes on other species of special concern (Lepidoptera: Saturniidae, Sphingiidae, Notodontidae, and Arctiidae). Ohio J. Sci. 90:33–40.
- OPLER, P. A. & G. O. KRIZEK. 1984. Butterflies east of the Great Plains. Johns Hopkins Univ. Press, Baltimore, Maryland. 294 pp.
- PANZER, R. D. STILLWAUGH, R. GNAEDINGER & G. DERKOVITZ. 1995. Prevalence of remnant dependence among the prairie- and savanna-inhabiting insects of the Chicago Region. Natural Areas Journal 15:101–116.
- RINGS, R. W., E. H. METZLER, F. J. ARNOLD & D. H. HARRIS. 1992. The owlet moths of Ohio: Order Lepidoptera, Family Noctuidae. Ohio Biol. Surv. Bull. New Series Vol. 9, No. 2. 219 pp.
- SHUEY, J. A., J. V. CALHOUN & D. C. IFTNER. 1987. Butterflies that are endangered, threatened, and of special concern in Ohio. Ohio J. Sci. 87:98–106.
- SHUEY, J. A., E. H. METZLER, D. C. IFTNER, J. V. CALHOUN, J. W. PEACOCK, R. A. WATKINS, J. D. HOOPER & W. F. BABCOCK. 1987. Status and habitats of potentially endangered Lepidoptera in Ohio. J. Lepid. Soc. 41:1–12.
- SWENGEL, A. B. 1994. Observations on the effects of fire on Karner blue butterflies, pp. 81–86. *In* D. A. Andow, R. J. Baker & C. P. Lane (eds.), Karner blue butterfly: symbol of a vanishing landscape. Misc. Publ. 84-1994, Minn. Agric. Exp. Stat., Univ. Minn. St. Paul.
- THOMAS, C. D. & S. HARRISON. 1992. Spatial dynamics of a patchily distributed butterfly species. J. Animal Ecol. 61:437–446.
- ZAREMBA, R. E. & M. PICKERING. 1994. Lupine ecology and management in New York State, pp. 201–208. *In* D. A. Andow, R. J. Baker & C. P. Lane (eds.), Karner blue butterfly: symbol of a vanishing landscape. Misc. Publ. 84-1994, Minn. Agric. Exp. Stat., Univ. Minn. St. Paul.

JOHN A. SHUEY, *The Nature Conservancy, Indiana Field Office, 1330 West 38th Street, Indianapolis, Indiana 46208, USA.*

Received for publication 22 August 1995; revised and accepted 13 July 1996.

Journal of the Lepidopterists' Society
51(3), 1997, 269–271

LIFE HISTORY NOTES FOR THE PALLID EMPEROR MOTH, *CIRINA FORDA* (SATURNIIDAE) IN NIGERIA

Additional key words: phenology, hostplants, Africa.

Cirina forda Westwood has long been known as a serious pest of the sheanut tree, *Vitellaria paradoxa* (Sapotaceae) in Nigeria (Golding 1929). Packard (1914) described the larva, and Boorman (1970) and Leleup and Beams (1969) provided brief accounts of the biology and phenology of this moth. Leleup and Deans (1969) reported *Erythropheum africanum* as a larval host in northern Zaire, but that the tree does not occur in Nigeria. The dried larvae of *C. forda* are referred to locally as “manimani,” and are of economic





FIG. 1. Phenology of *Cirina forda* in Nigeria, 1988-1991.

importance as a food item among the Nupe tribe of Nigeria. For many decades, wild larvae have been collected, dried and sold to neighboring states within and outside the country, and the insect is considered a delicacy (Ande 1991, Fasoranti & Ajiboye 1993). We report here on the life history and biology of *C. forda* in Nigeria, and its association with *Vittelaria paradoxa*.

About 900 late instar *C. forda* were collected from *V. paradoxa* plants in August 1988, between km 107 and 108 on the Mokwa-Bida road in Niger State, Nigeria (9°05'N, 5°59'E). Subsequently, rearings were conducted in the laboratory at the University of Ilorin between 1989 and 1991. Egg clusters laid in the laboratory were observed daily for hatching. Larvae from the same egg clutch were reared separately on cut and potted twigs of *V. paradoxa* and again on *V. paradoxa* plants outside in an ornamental garden. Twigs were replaced as necessary to maintain freshness, and larvae were transferred carefully by hand and placed on the mid rib or leaf margin of new foliage. A wooden cage (30 cm × 30 cm × 50 cm) with ten compartments was filled to a depth of 20 cm with soil. Ten larvae were then placed in each compartment. As soon as pupation began, the soil was in all compartments examined for pre-pupae and pupae. Pre-pupa duration was defined as the pe-



FIGS. 2-6. 2, egg cluster. 3, pre-pupa. 4, fully formed pupa. 5, late instar. 6, adults (female above, male below).

riod between soil penetration and actual pupation (in days). Subsequently, 30 pre-pupae were randomly selected and placed individually in soil 10 cm deep, each in cylindrical paper eclosion chambers (8 cm diam \times 15 cm deep \times 2 cm thick). The open end of the cylinder was covered with a nylon mesh secured by a rubber band. Each compartment was examined daily for emerging adults. Pupal duration was calculated as the mean number of days between pupal formation and date of adult emergence. Notes were also kept on adult longevity.

Fig. 1 shows the phenology of *C. forda* for the period between 1988 and 1991. Adult moths lived for between 36 and 48 hours (mean = 39.7) and were found primarily in May, with peak oviposition at the end of the month. Figs. 2–6 show the immature stages of *C. forda*. The egg (Fig. 2) hatches after an incubation period of 30 to 34 days (mean = 31.8) into an active and voraciously feeding larva, and passes through 5 to 6 instars (Fig. 5) in 42 to 50 days (mean = 47.5) between June and August. By the first few days in August, most of the larvae reach the pre-pupal stage (Fig. 3) and burrow into the soil. The pre-pupa develops into a pupa (Fig. 4) in 6–7 days and remains in diapause for 9 months (261 to 296 days, mean = 267.5). Adult moths (Fig. 6) emerge in May of the following year.

The life cycle of *C. forda* is tightly linked to the biology of its host, *Vittelaria paradoxa*. The only savannah species of the family Sapotaceae in Nigeria, *V. paradoxa* blossoms fully between May and August when mature fruits become available, and sheds leaves between November and February (Kaay et al. 1964). May and August is when the majority of *C. forda* larvae are developing in the field, and pupation takes place during the dry months of November and April. *C. forda* is univoltine in Nigeria and the phenology reported here agrees with those given by Golding (1929) and Boorman (1978). However, Leleup and Deams (1969) indicate that the active period for *C. forda* in Zaire is between June and September during dry months. In Nigeria, the active periods occur during the wet months of May and August.

ACKNOWLEDGMENTS

We acknowledge a postgraduate fellowship and Senate Research Grant awarded by the University of Ilorin, Ilorin, Nigeria.

LITERATURE CITED

- ANDE, A. T. 1991. Some aspects of the biology of *Cirina forda* Westwood (Lepidoptera: Saturniidae) Unpubl. Ph. D. Thesis, Dept. Biol. Sciences, Univ. Ilorin, Nigeria. 327 pp.
- BOORMAN, J. P. T. 1970. The emperor moths (Saturniidae) of Nigeria. *Nigerian Field* 35:99–122.
- . 1978. *West African Butterflies and moths*. Longman Group Ltd., London. 2nd Ed. 79 pp.
- FASORANTI, J. O. & D. O. AJIBOYE. 1993. Some edible insects of Kwara State, Nigeria. *Amer. Entomol.* 93:113–116.
- GOLDING, F. D. 1929. Preliminary notes on the pests of sheanut tree in Northern Nigeria. *Bull. Dept. Agric. Nigeria* 8:101–103.
- KEAY, R. W. J., ONOCHIE, C. F. A. & D. P. STANDFIELD. 1964. *Nigerian trees*. Dept. Forest Research. Ibadan, Nigeria. 495 pp.
- LELEUP, N & H. DAEMS. 1969. Les chenilles alimentaires du kwango causes de leur rarefaction et mesures pre-consees pour y remedier. *J. Agric. Trop. & Botany Appl.* 16:1–21.

A. TAIWO ANDE AND J. OLANIYAN FASORANTI *Department of Biological Sciences, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria.*

Received for publication 6 December 1994; revised and accepted 26 January 1997.